

EVERYONE WILL NOTICE. *No One Will Know.*

Get natural-looking results that last
for cheeks, lines, and lips.



Actual patient treated with JUVÉDERM VOLUMA® XC, JUVÉDERM VOLLURE™ XC, and JUVÉDERM VOLBELLA® XC injectable gels. Results may vary.



**Actual patient.
Results may vary.**

Unretouched photos taken before treatment and 1 month after treatment. A total of 1.4 mL of JUVÉDERM VOLUMA® XC was injected into the cheek area. A total of 2.9 mL of JUVÉDERM VOLLURE™ XC was injected into the parentheses and marionette lines. A total of 2.2 mL of JUVÉDERM VOLBELLA® XC was injected into the vertical lip lines and lips.

JUVÉDERM® offers a wide range of treatment options that can be customized to your individual needs.
Ask your aesthetic specialist which treatments may be right for you.

JUVÉDERM® Injectable Gel Fillers Important Information

APPROVED USES

JUVÉDERM VOLUMA® XC injectable gel is for deep injection in the cheek area to correct age-related volume loss in adults over 21.

JUVÉDERM® XC and JUVÉDERM VOLLURE™ XC injectable gels are for injection into the facial tissue for the correction of moderate to severe facial wrinkles and folds, such as nasolabial folds.

JUVÉDERM® Ultra XC is for injection into the lips and perioral area for lip augmentation in adults over the age of 21.

JUVÉDERM VOLBELLA® XC injectable gel is for injection into the lips for lip augmentation and for correction of perioral lines in adults over the age of 21.

IMPORTANT SAFETY INFORMATION

Are there any reasons why I should not receive any JUVÉDERM® injectable gel formulation?

Do not use these products if you have a history of multiple severe allergies or severe allergic reactions (anaphylaxis), or if you are allergic to lidocaine or the Gram-positive bacterial proteins used in these products.

What precautions should my doctor advise me about?

- Tell your doctor if you are pregnant or breastfeeding. The safety of these products for use during pregnancy or while breastfeeding has not been studied

- The safety of JUVÉDERM VOLUMA® XC in patients under 35 years or over 65 years, the safety of JUVÉDERM® XC and JUVÉDERM® Ultra XC injectable gels in patients under 18 years, and the safety of JUVÉDERM VOLLURE™ XC and JUVÉDERM VOLBELLA® XC in patients under 22 years has not been studied
- The safety and effectiveness of JUVÉDERM VOLUMA® XC in areas other than the cheek area, JUVÉDERM® XC and JUVÉDERM VOLLURE™ XC for areas other than facial wrinkles and folds, and JUVÉDERM® Ultra XC and JUVÉDERM VOLBELLA® XC in areas other than the lips and perioral area have not been established in clinical studies
- Tell your doctor if you have a history of excessive scarring (eg, hypertrophic scarring and keloid formations) or pigmentation disorders, as use of these products may result in additional scars or changes in pigmentation
- Tell your doctor if you are planning other laser treatments or a chemical peel, as there is a possible risk of inflammation at the treatment site if these procedures are performed after treatment
- Patients who experience skin injury near the site of injection with these products may be at a higher risk for side effects
- Tell your doctor if you are on immunosuppressive therapy used to decrease the body's immune response, as use of these products may result in an increased risk of infection
- Tell your doctor if you are using medications that can prolong bleeding, such as aspirin, ibuprofen, or other blood thinners, as this may result in increased bruising or bleeding at the injection site
- Minimize strenuous exercise, exposure to extensive sun or heat, and alcoholic beverages within the first 24 hours following treatment



© 2017 Allergan. All rights reserved. All trademarks are the property of their respective owners.
Juvéderm.com Re-order: JUV106073 03/17 170190

What are possible side effects?

The most commonly reported side effects with JUVÉDERM® injectable gels included injection-site redness, swelling, pain, tenderness, firmness, lumps/bumps, bruising, discoloration, and itching. For JUVÉDERM VOLBELLA® XC, dryness was also reported. For JUVÉDERM VOLUMA® XC, most side effects were moderate and lasted 2 to 4 weeks. For JUVÉDERM® XC, JUVÉDERM VOLLURE™ XC, and JUVÉDERM® Ultra XC injectable gels, most side effects were mild or moderate and lasted 14 days or less. For JUVÉDERM VOLBELLA® XC, most side effects were mild or moderate and lasted 30 days or less.

One of the risks with using these products is unintentional injection into a blood vessel, and, while rare, the complications can be serious and may be permanent. These complications, which have been reported for facial injections, can include vision abnormalities, blindness, stroke, temporary scabs, or permanent scarring.

As with all skin injection procedures, there is a risk of infection.

To report a side effect with any JUVÉDERM® product, please call Allergan at 1-800-433-8871. Please visit Juvéderm.com or talk to your doctor for more information.

Available by prescription only.

